

Histone deacetylase 8 Polyclonal Conjugated Antibody

Catalog No: #C42203

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Package Size: #C42203-AF350 100ul #C42203-AF405 100ul #C42203-AF488 100ul

#C42203-AF555 100ul #C42203-AF594 100ul #C42203-AF647 100ul

#C42203-AF680 100ul #C42203-AF750 100ul #C42203-Biotin 100ul

Description

Product Name	Histone deacetylase 8 Polyclonal Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of total Histone deacetylase 8 polyclonal antibody.
Immunogen Description	Recombinant human Histone deacetylase 8 protein $\Omega\frac{1}{2}\Omega\frac{1}{2}$ -377aa $\Omega\frac{1}{2}\Omega\frac{1}{2}$
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	HDAC8,HDACL1,CDA07
Accession No.	Swiss-Prot#:Q9BY41
Uniprot	Q9BY41
GeneID	55869;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Also involved in the deacetylation of cohesin complex protein SMC3 regulating release of cohesin complexes from chromatin. May play a role in smooth muscle cell contractility.

Note: This product is for in vitro research use only