

Lysosomal acid lipase/cholesteryl ester hydrolase Polyclonal Conjugated Antibody

Catalog No: #C42238

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Package Size: #C42238-AF350 100ul #C42238-AF405 100ul #C42238-AF488 100ul

#C42238-AF555 100ul #C42238-AF594 100ul #C42238-AF647 100ul

#C42238-AF680 100ul #C42238-AF750 100ul #C42238-Biotin 100ul

Description

| | |
|-----------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Name | Lysosomal acid lipase/cholesteryl ester hydrolase Polyclonal Conjugated Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous level of total Lysosomal acid lipase/cholesteryl ester hydrolase polyclonal antibody. |
| Immunogen Description | Recombinant human Lysosomal acid lipase/cholesteryl ester hydrolase protein |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | Cholesteryl esterase Lipase A Sterol esterase LIPA |
| Accession No. | Swiss-Prot#:P38571 |
| Uniprot | P38571 |
| GeneID | 3988; |
| Excitation Emission | AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm |
| Calculated MW | 42 |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Crucial for the intracellular hydrolysis of cholesteryl esters and triglycerides that have been internalized via receptor-mediated endocytosis of lipoprotein particles. Important in mediating the effect of LDL (low density lipoprotein) uptake on suppression of hydroxymethylglutaryl-CoA reductase and activation of endogenous cellular cholesteryl ester formation.

Note: This product is for in vitro research use only