

## Prefoldin subunit 5 Polyclonal Conjugated Antibody

Catalog No: #C42255



Package Size: #C42255-AF350 100ul #C42255-AF405 100ul #C42255-AF488 100ul  
 #C42255-AF555 100ul #C42255-AF594 100ul #C42255-AF647 100ul  
 #C42255-AF680 100ul #C42255-AF750 100ul #C42255-Biotin 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)  
 Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Prefoldin subunit 5 Polyclonal Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of total Prefoldin subunit 5 polyclonal antibody.
Immunogen Description	Recombinant human Prefoldin subunit 5 protein
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	PFDN5
Accession No.	Swiss-Prot#:Q99471
Uniprot	Q99471
GeneID	5204;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

## Application Details

## Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Background

---

Binds specifically to cytosolic chaperonin (c-CPN) and transfers target proteins to it. Binds to nascent polypeptide chain and promotes folding in an environment in which there are many competing pathways for nonnative proteins. Heterohexamer of two PFD-alpha type and four PFD-beta type subunits. Binds to MYC; interacts with its N-terminal domain. Represses the transcriptional activity of MYC. Highly expressed in pancreas and skeletal muscle and moderately in other tissues. Belongs to the prefoldin subunit alpha family.

---

Note: This product is for in vitro research use only