

Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit DAD1 Polyclonal Conjugated Antibody

Catalog No: #C42387

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Package Size: #C42387-AF350 100ul #C42387-AF405 100ul #C42387-AF488 100ul

#C42387-AF555 100ul #C42387-AF594 100ul #C42387-AF647 100ul

#C42387-AF680 100ul #C42387-AF750 100ul #C42387-Biotin 100ul

Description

Product Name	Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit DAD1 Polyclonal Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of total Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit DAD1 polyclonal antibody.
Immunogen Description	Recombinant human Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit DAD1 protein
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	Defender against cell death 1
Accession No.	Swiss-Prot#:P61803
Uniprot	P61803
GeneID	1603;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	12 .4
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Component of the N-oligosaccharyl transferase enzyme which catalyzes the transfer of a high mannose oligosaccharide from a lipid-linked oligosaccharide donor to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains. N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). Loss of the DAD1 protein triggers apoptosis.

Note: This product is for in vitro research use only