

Tumor necrosis factor receptor superfamily member 6 Polyclonal Conjugated Antibody



Catalog No: #C42455

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Package Size: #C42455-AF350 100ul #C42455-AF405 100ul #C42455-AF488 100ul

#C42455-AF555 100ul #C42455-AF594 100ul #C42455-AF647 100ul

#C42455-AF680 100ul #C42455-AF750 100ul #C42455-Biotin 100ul

Description

Product Name	Tumor necrosis factor receptor superfamily member 6 Polyclonal Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of total Tumor necrosis factor receptor superfamily member 6 polyclonal antibody.
Immunogen Description	Recombinant human Tumor necrosis factor receptor superfamily member 6 protein
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	Apo-1 antigen Apoptosis-mediating surface antigen FAS FASLG receptor CD_antigen=CD95
Accession No.	Swiss-Prot#:P25445
Uniprot	P25445
GeneID	355;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	36.8
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Receptor for TNFSF6/FASLG. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. FAS-mediated apoptosis may have a role in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T-cells, or both. The secreted isoforms 2 to 6 block apoptosis (in vitro).

Note: This product is for in vitro research use only