

Amyloid beta A4 protein Polyclonal Conjugated Antibody



Catalog No: #C42478

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Package Size: #C42478-AF350 100ul #C42478-AF405 100ul #C42478-AF488 100ul

#C42478-AF555 100ul #C42478-AF594 100ul #C42478-AF647 100ul

#C42478-AF680 100ul #C42478-AF750 100ul #C42478-Biotin 100ul

Description

Product Name	Amyloid beta A4 protein Polyclonal Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of total Amyloid beta A4 protein polyclonal antibody.
Immunogen Description	Recombinant human Amyloid beta A4 protein
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	ABPP APPI APP Alzheimer disease amyloid protein Cerebral vascular amyloid peptide CVAP PreA4 Protease nexin-II PN-II
Accession No.	Swiss-Prot#:P05067
Uniprot	P05067
GeneID	351;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Background

Functions as a cell surface receptor and performs physiological functions on the surface of neurons relevant to neurite growth, neuronal adhesion and axonogenesis. Involved in cell mobility and transcription regulation through protein-protein interactions. Can promote transcription activation through binding to APBB1-KAT5 and inhibits Notch signaling through interaction with Numb. Couples to apoptosis-inducing pathways such as those mediated by G(O) and JIP. Inhibits G(o) alpha ATPase activity By similarity. Acts as a kinesin I membrane receptor, mediating the axonal transport of beta-secretase and presenilin 1. Involved in copper homeostasis/oxidative stress through copper ion reduction. In vitro, copper-metallated APP induces neuronal death directly or is potentiated through Cu²⁺-mediated low-density lipoprotein oxidation. Can regulate neurite outgrowth through binding to components of the extracellular matrix such as heparin and collagen I and IV. The splice isoforms that contain the BPTI domain possess protease inhibitor activity. Induces a AGER-dependent pathway that involves activation of p38 MAPK, resulting in internalization of amyloid-beta peptide and leading to mitochondrial dysfunction in cultured cortical neurons. Ref.36 Ref.63 Ref.65 Ref.88 Ref.89 Beta-amyloid peptides are lipophilic metal chelators with metal-reducing activity. Bind transient metals such as copper, zinc and iron. In vitro, can reduce Cu²⁺ and Fe³⁺ to Cu⁺ and Fe²⁺, respectively. Beta-amyloid 42 is a more effective reductant than beta-amyloid 40. Beta-amyloid peptides bind to lipoproteins and apolipoproteins E and J in the CSF and to HDL particles in plasma, inhibiting metal-catalyzed oxidation of lipoproteins. Beta-APP42 may activate mononuclear phagocytes in the brain and elicit inflammatory responses. Promotes both tau aggregation and TPK II-mediated phosphorylation. Interaction with overexpressed HADH2 leads to oxidative stress and neurotoxicity. Ref.36 Ref.63 Ref.65 Ref.88 Ref.89 Appicans elicit adhesion of neural cells to the extracellular matrix and may regulate neurite outgrowth in the brain By similarity. Ref.36 Ref.63 Ref.65 Ref.88 Ref.89 The gamma-CTF peptides as well as the caspase-cleaved peptides, including C31, are potent enhancers of neuronal apoptosis. Ref.36 Ref.63 Ref.65 Ref.88 Ref.89 N-APP binds TNFRSF21 triggering caspase activation and degeneration of both neuronal cell bodies (via caspase-3) and axons (via caspase-6). Ref.36 Ref.63 Ref.65 Ref.88 Ref.89

Note: This product is for in vitro research use only