## CDH16 Conjugated Antibody

Catalog No: #C46449



 Package Size:
 #C46449-AF350 100ul
 #C46449-AF405 100ul
 #C46449-AF488 100ul

 #C46449-AF555 100ul
 #C46449-AF594 100ul
 #C46449-AF647 100ul

 #C46449-AF680 100ul
 #C46449-AF750 100ul
 #C46449-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

## Description

Product Name	CDH16 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous levels of total CDH16 protein.
Immunogen Description	Synthetic protein corresponding to residues near the C terminal of human CDH16
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Accession No.	Swiss-Prot#:O75309NCBI Gene ID:1014NCBI Protein#:BC027912
Uniprot	O75309
GenelD	1014;
Excitation Emission	AF350: 346nm/442nm
	AF405: 401nm/421nm
	AF488: 493nm/519nm
	AF555: 555nm/565nm
	AF594: 591nm/614nm
	AF647: 651nm/667nm
	AF680: 679nm/702nm
	AF750: 749nm/775nm
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application De	tails
Suggested Dilution	:
AF350 conjugated:	most applic
AF405 conjugated:	most applic
AF488 conjugated:	most applic
AF555 conjugated:	most applic
AF594 conjugated:	most applic
AF647 conjugated:	most applic
AF680 conjugated:	most applic
AF750 conjugated:	most applic
Biotin conjugated: \	working with

## Background

This gene is a member of the cadherin superfamily, genes encoding calcium-dependent, membrane-associated glycoproteins. Mapped to a previously identified cluster of cadherin genes on chromosome 16q22.1, the gene localizes with superfamily members CDH1, CDH3, CDH5, CDH8 and CDH11. The protein consists of an extracellular domain containing 6 cadherin domains, a transmembrane region and a truncated cytoplasmic domain but lacks the prosequence and tripeptide HAV adhesion recognition sequence typical of most classical cadherins. Expression is exclusively in kidney, where the protein functions as the principal mediator of homotypic cellular recognition, playing a role in the morphogenic direction of tissue development. Alternatively spliced transcript variants encoding distinct isoforms have been identified.?

Note: This product is for in vitro research use only