

## CCL11 Conjugated Antibody

Catalog No: #C46940



Package Size: #C46940-AF350 100ul #C46940-AF405 100ul #C46940-AF488 100ul  
 #C46940-AF555 100ul #C46940-AF594 100ul #C46940-AF647 100ul  
 #C46940-AF680 100ul #C46940-AF750 100ul #C46940-Biotin 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)  
 Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

|                       |  |
|-----------------------|--|
| Product Name          | CCL11 Conjugated Antibody  |
| Host Species          | Rabbit   |
| Clonality             | Polyclonal   |
| Species Reactivity    | Hu   |
| Specificity           | The antibody detects endogenous levels of total CCL11 protein.   |
| Immunogen Description | Synthetic peptide of human CCL11   |
| Conjugates            | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750   |
| Other Names           | SCYA11   |
| Accession No.         | Swiss-Prot#:P51671 NCBI Gene ID:6356NCBI Protein#:NP_002977  |
| Uniprot               | P51671   |
| GeneID                | 6356;  |
| Excitation Emission   | AF350: 346nm/442nm<br>AF405: 401nm/421nm<br>AF488: 493nm/519nm<br>AF555: 555nm/565nm<br>AF594: 591nm/614nm<br>AF647: 651nm/667nm<br>AF680: 679nm/702nm<br>AF750: 749nm/775nm |
| Formulation           | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide  |
| Storage               | Store at 4°C in dark for 6 months  |

## Application Details

## Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Background

---

This antimicrobial gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of the N-terminal cysteine residues of the mature peptide. This chemokine, a member of the CC subfamily, displays chemotactic activity for eosinophils, but not mononuclear cells or neutrophils. This eosinophil-specific chemokine is thought to be involved in eosinophilic inflammatory diseases such as atopic dermatitis, allergic rhinitis, asthma and parasitic infections.?

---

Note: This product is for in vitro research use only