

DAGLB Conjugated Antibody

Catalog No: #C47015



Package Size: #C47015-AF350 100ul #C47015-AF405 100ul #C47015-AF488 100ul
 #C47015-AF555 100ul #C47015-AF594 100ul #C47015-AF647 100ul
 #C47015-AF680 100ul #C47015-AF750 100ul #C47015-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	DAGLB Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous levels of total DAGLB protein.
Immunogen Description	Synthetic peptide of human DAGLB
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	KCCR13L; DAGLBETA
Accession No.	Swiss-Prot#:Q8NCG7 NCBI Gene ID:221955NCBI Protein#:NP_631918
Uniprot	Q8NCG7
GeneID	221955;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Members of the AB hydrolase superfamily have diverse catalytic functions and play a crucial role in the metabolism of lipids. DAGL β (diacylglycerol lipase beta), also known as KCCR13L, is a 672 amino acid multi-pass membrane protein that belongs to the AB hydrolase superfamily. DAGL β uses calcium as a cofactor to catalyze the hydrolysis of diacylglycerol (DAG) to 2-arachidonoyl-glycerol (2-AG), a reaction that is required for axonal growth and for retrograde synaptic signaling at mature synapses. DAGL β functions at an optimal pH of 7 and its activity is inhibited by p-hydroxy-mercuri-benzoate and HgCl₂, but not PMSF. There are three isoforms of DAGL β that are produced as a result of alternative splicing events.

Note: This product is for in vitro research use only