DBX2 Conjugated Antibody

Catalog No: #C47023

SAB Signalway Antibody

Package Size: #C47023-AF350 100ul #C47023-AF405 100ul #C47023-AF488 100ul

#C47023-AF555 100ul #C47023-AF594 100ul #C47023-AF647 100ul

#C47023-AF680 100ul #C47023-AF750 100ul #C47023-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

| Product Name | DBX2 Conjugated Antibody |
|-----------------------|---|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous levels of total DBX2 protein. |
| Immunogen Description | Synthetic peptide of human DBX2 |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Accession No. | Swiss-Prot#:Q6ZNG2 NCBI Gene ID:440097NCBI Protein#:NP_001004329 |
| Uniprot | Q6ZNG2 |
| GeneID | 440097; |
| Excitation Emission | AF350: 346nm/442nm |
| | AF405: 401nm/421nm |
| | AF488: 493nm/519nm |
| | AF555: 555nm/565nm |
| | AF594: 591nm/614nm |
| | AF647: 651nm/667nm |
| | AF680: 679nm/702nm |
| | AF750: 749nm/775nm |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
AF405 conjugated: most applications: 1: 50 - 1: 250
AF488 conjugated: most applications: 1: 50 - 1: 250
AF555 conjugated: most applications: 1: 50 - 1: 250
AF594 conjugated: most applications: 1: 50 - 1: 250
AF647 conjugated: most applications: 1: 50 - 1: 250
AF680 conjugated: most applications: 1: 50 - 1: 250
AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

DBX2 is a member of the developing brain homeobox (DBX) protein family, but while the related protein DBX1 is expressed in various regions of the developing brain, DBX2 shows a more restricted pattern of expression in the brain, and is also expressed in some mesenchymal cells such as limb buds and tooth germs. It is thought that DBX1 and DBX2 promote the development of a subset of interneurons, some of which help mediate left-right coordination of locomotor activity. In Xenopus, DBX2 is involved in primary neurogenesis and early neural plate patterning, and is thought to act as a cross-repressive partner of NKX6-2 in the patterning of the ventral neural tube.

Note: This product is for in vitro research use only