

## DCTD Conjugated Antibody

Catalog No: #C47030



Package Size: #C47030-AF350 100ul #C47030-AF405 100ul #C47030-AF488 100ul  
 #C47030-AF555 100ul #C47030-AF594 100ul #C47030-AF647 100ul  
 #C47030-AF680 100ul #C47030-AF750 100ul #C47030-Biotin 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)  
 Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

|                       |  |
|-----------------------|--|
| Product Name          | DCTD Conjugated Antibody   |
| Host Species          | Rabbit   |
| Clonality             | Polyclonal   |
| Species Reactivity    | Hu   |
| Specificity           | The antibody detects endogenous levels of total DCTD protein.  |
| Immunogen Description | Full length fusion protein   |
| Conjugates            | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750   |
| Accession No.         | Swiss-Prot#:P32321NCBI Gene ID:1635NCBI Protein#:BC001286  |
| Uniprot               | P32321   |
| GeneID                | 1635;  |
| Excitation Emission   | AF350: 346nm/442nm<br>AF405: 401nm/421nm<br>AF488: 493nm/519nm<br>AF555: 555nm/565nm<br>AF594: 591nm/614nm<br>AF647: 651nm/667nm<br>AF680: 679nm/702nm<br>AF750: 749nm/775nm |
| Formulation           | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide  |
| Storage               | Store at 4°C in dark for 6 months  |

## Application Details

## Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Background

---

The protein encoded by this gene catalyzes the deamination of dCMP to dUMP, the nucleotide substrate for thymidylate synthase. The encoded protein is allosterically activated by dCTP and inhibited by dTTP, and is found as a homohexamer. This protein uses zinc as a cofactor for its activity. Two transcript variants encoding different isoforms have been found for this gene.

---

Note: This product is for in vitro research use only