

Hsc70 Conjugated Antibody

Catalog No: #C48246



Package Size: #C48246-AF350 100ul #C48246-AF405 100ul #C48246-AF488 100ul
 #C48246-AF555 100ul #C48246-AF594 100ul #C48246-AF647 100ul
 #C48246-AF680 100ul #C48246-AF750 100ul #C48246-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	Hsc70 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu, Ms, Rt
Immunogen Description	Peptide
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	2410008N15Rik antibody Constitutive heat shock protein 70 antibody Epididymis luminal protein 33 antibody Epididymis secretory sperm binding protein Li 72p antibody Heat shock 70 kDa protein 8 antibody Heat shock 70kD protein 10 antibody Heat shock 70kD protein 8 antibody Heat shock 70kDa protein 8 antibody Heat shock cognate 71 kDa protein antibody Heat shock cognate protein 54 antibody Heat shock cognate protein 71 kDa antibody Heat shock protein 8 antibody Heat shock protein A8 antibody Heat shock protein family A (Hsp70) member 8 antibody Heat-shock70-KD protein 10, formerly antibody HEL 33 antibody HEL S 72p antibody HSC54 antibody HSC71 antibody Hsc73 antibody HSP71 antibody HSP73 antibody HSP7C_HUMAN antibody HSPA10 antibody HSPA8 antibody LAP1 antibody Lipopolysaccharide associated protein 1 antibody LPS associated protein 1 antibody LPS associated protein antibody MGC102007 antibody MGC106514 antibody MGC114311 antibody MGC118485 antibody MGC131511 antibody MGC29929 antibody N-myristoyltransferase inhibitor protein 71 antibody NIP71 antibody
Accession No.	Swiss-Prot#:P11142
Uniprot	P11142
GenelD	3312;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	71 kDa
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

The HSP 70 family is composed of four highly conserved proteins: HSP 70, HSC 70, GRP 75 and GRP 78. These proteins serve a variety of roles: they act as molecular chaperones facilitating the assembly of multi-protein complexes, participate in the translocation of polypeptides across cell membranes and to the nucleus, and aid in the proper folding of nascent polypeptide chains. All members of the family, except HSP 70, are constitutively expressed in primate cells. HSP 70 expression is strongly induced in response to heat stress. HSP 70 and HSC 70 play key roles in the cytosolic endoplasmic reticulum and mitochondrial import machinery and are found in both the cytosol and nucleus of mammalian cells. Both HSP 70 and HSC 70 are involved in the chaperoning of nascent polypeptide chains and in protecting cells against the accumulation of improperly folded proteins. GRP 78 is localized in the endoplasmic reticulum, where it receives imported secretory proteins and is involved in the folding and translocation of nascent peptide chains. GRP 75 expression is restricted to the mitochondrial matrix and aids in the translocation and folding of nascent polypeptide chains of both nuclear and mitochondrial origin. GRP 75 and GRP 78 are unresponsive to heat stress and are induced by glucose deprivation. It has been postulated that members of the HSP 70 family act as force-generating motors, relying on the hydrolysis of ATP for their activity.

Note: This product is for in vitro research use only