

ANAPC11 Conjugated Antibody

Catalog No: #C48441



Package Size: #C48441-AF350 100ul #C48441-AF405 100ul #C48441-AF488 100ul
 #C48441-AF555 100ul #C48441-AF594 100ul #C48441-AF647 100ul
 #C48441-AF680 100ul #C48441-AF750 100ul #C48441-Biotin 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	ANAPC11 Conjugated Antibody
Host Species	Mouse
Clonality	Monoclonal
Species Reactivity	Hu
Immunogen Description	Recombinant protein
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	ANAPC 11 antibody ANAPC11 antibody Anaphase promoting complex subunit 11 (yeast APC11 homolog) antibody Anaphase promoting complex subunit 11 antibody Anaphase promoting complex subunit 11 homolog (yeast) antibody Anaphase promoting complex subunit 11 homolog antibody Anaphase-promoting complex subunit 11 antibody Apc 11 antibody Apc 11p antibody APC11 anaphase promoting complex subunit 11 homolog (yeast) antibody APC11 anaphase promoting complex subunit 11 homolog antibody APC11 antibody APC11_HUMAN antibody Apc11p antibody Cyclosome subunit 11 antibody Hepatocellular carcinoma associated RING finger protein antibody Hepatocellular carcinoma-associated RING finger protein antibody HSPC 214 antibody HSPC214 antibody MGC882 antibody Yeast APC 11 homolog antibody Yeast APC11 homolog antibody
Accession No.	Swiss-Prot#:Q9NYG5
Uniprot	Q9NYG5
GenElD	51529;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	10 kDa
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Comprising more than ten subunits, the anaphase-promoting complex (APC) acts in a cell-cycle dependent manner to promote the separation of sister chromatids during the transition between metaphase and anaphase in mitosis. APC, or cyclosome, accomplishes this progression through the ubiquitination of mitotic cyclins and other regulatory proteins that are targeted for destruction during cell division. APC is phosphorylated, and thus activated, by protein kinases Cdk1/cyclin B and polo-like kinase (Plk). APC is under tight control by a number of regulatory factors, including CDC20, CDH1 and MAD2. Specifically, CDC20 and CDH1 directly bind to APC and activates APC's cyclin-ubiquitination activity. In contrast, MAD2 inhibits APC by forming a ternary complex with CDC20 and APC; thus preventing APC activation. APC11 is a RING-H2 finger protein that allows for the synthesis of multiubiquitin chains in the presence of Ubiquitin carrier protein 4 (Ubc4) and ubiquitin conjugating enzyme (E2). In addition, a heterodimeric complex of either Ubc4 or UbcH10 with APC11 and APC2 catalyzes the ubiquitination of human securin and cyclin B1.

Note: This product is for in vitro research use only