

MCSF Conjugated Antibody

Catalog No: #C48850



Package Size: #C48850-AF350 100ul #C48850-AF405 100ul #C48850-AF488 100ul
 #C48850-AF555 100ul #C48850-AF594 100ul #C48850-AF647 100ul
 #C48850-AF680 100ul #C48850-AF750 100ul #C48850-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

| | |
|-----------------------|---|
| Product Name | MCSF Conjugated Antibody |
| Host Species | Rabbit |
| Clonality | Monoclonal |
| Species Reactivity | Hu |
| Immunogen Description | recombinant protein |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | Colony stimulating factor 1 (macrophage) antibody Colony stimulating factor 1 antibody Colony stimulating factor macrophage specific antibody CSF 1 antibody CSF-1 antibody CSF1 antibody CSF1_HUMAN antibody Csfm antibody Lanimostim antibody M-CSF antibody M-CSF antibody Macrophage Colony Stimulating Factor 1 antibody Macrophage colony stimulating factor antibody MCSF antibody MGC31930 antibody Processed macrophage colony-stimulating factor 1 antibody |
| Accession No. | Swiss-Prot#:P09603 |
| Uniprot | P09603 |
| GeneID | 1435; |
| Excitation Emission | AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm |
| Calculated MW | 60 kDa |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

The macrophage colony-stimulating factor (M-CSF), also designated CSF-1, was originally discovered in serum, urine and other biological fluids as a factor that can stimulate the formation of macrophage colonies from bone marrow hematopoietic progenitor cells. M-CSF is a homodimeric cytokine that is produced by fibroblasts, epithelial cells, bone marrow stromal cells, osteoblasts, keratinocytes, macrophages, T cells and B cells. M-CSF is a glycoprotein required for the proliferation and differentiation of mononuclear phagocytes, including osteoclasts. M-CSF has also been identified as an important mediator of the inflammatory response and can regulate the release of proinflammatory cytokines from macrophages. M-CSF exerts its pleiotropic effects by binding to a single type of high affinity cell surface receptor that is encoded by the c-Fms proto-oncogene.

Note: This product is for in vitro research use only