

Cdk8 Conjugated Antibody

Catalog No: #C49116



Package Size: #C49116-AF350 100ul #C49116-AF405 100ul #C49116-AF488 100ul
 #C49116-AF555 100ul #C49116-AF594 100ul #C49116-AF647 100ul
 #C49116-AF680 100ul #C49116-AF750 100ul #C49116-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	Cdk8 Conjugated Antibody
Host Species	Rabbit
Clonality	Monoclonal
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	cdk8 antibody CDK8 protein kinase antibody CDK8_HUMAN antibody Cell Division Kinase 8 antibody Cell division protein kinase 8 antibody Cyclin Dependent kinase 8 antibody Cyclin-dependent kinase 8 antibody K35 antibody Mediator complex subunit cdk8 antibody Mediator of RNA polymerase II transcription subunit cdk8 antibody Protein kinase K35 antibody
Accession No.	Swiss-Prot#:P49336
Uniprot	P49336
GeneID	1024;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	53 kDa
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Cell cycle progression is controlled in part by a family of cyclin proteins and cyclin dependent kinases (Cdks). Cdk proteins work in concert with the cyclins to phosphorylate key substrates involved in each phase of cell cycle progression. Another family of proteins, Cdk inhibitors, also plays a role in regulating cell cycle by binding to cyclin-Cdk complexes and modulating their activity. Several Cdk proteins have been identified, including Cdk2-Cdk8, PCTAIRE-1 α Ω $\frac{1}{2}$ C3, PITALRE and PITSLRE. Large complexes containing Cdk8, cyclin C and the large subunit of RNA polymerase II have been identified. Cdk8 is thought to regulate RNA polymerase II function in conjunction with cyclin C. Cdk8 has been demonstrated to function as a transcriptional activator when fused to the DNA binding domain of GAL4.

Note: This product is for in vitro research use only