Product Datasheet

Glutamate receptor 1 Conjugated Antibody

Catalog No: #C49118



Package Size: #C49118-AF350 100ul #C49118-AF405 100ul #C49118-AF488 100ul #C49118-AF555 100ul #C49118-AF594 100ul #C49118-AF647 100ul #C49118-AF680 100ul #C49118-AF750 100ul #C49118-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	Glutamate receptor 1 Conjugated Antibody		
Host Species	Rabbit		
Clonality	Monoclonal		
Species Reactivity	Hu, Ms, Rt		
Immunogen Description	recombinant protein		
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750		
Other Names	GLUR 1 antibody GLUR A antibody AMPA 1 antibody AMPA selective glutamate receptor 1 antibody		
	AMPA-selective glutamate receptor 1 antibody GluA1 antibody GLUH1 antibody GluR K1 antibody GluR-1		
	antibody GluR-A antibody GluR-K1 antibody GLUR1 antibody GLURA antibody GluRK1 antibody		
	Glutamate receptor 1 antibody Glutamate receptor ionotropic AMPA 1 antibody Glutamate receptor		
	ionotropic antibody Glutamate receptor, ionotropic, AMPA 1 antibody Gria1 antibody GRIA1_HUMAN		
	antibody HBGR1 antibody MGC133252 antibody OTTHUMP00000160643 antibody		
	OTTHUMP00000165781 antibody OTTHUMP00000224241 antibody OTTHUMP00000224242 antibody		
	OTTHUMP00000224243 antibody		
Accession No.	Swiss-Prot#:P42261		
Accession No. Uniprot	Swiss-Prot#:P42261 P42261		
Accession No. Uniprot GeneID	Swiss-Prot#:P42261 P42261 2890;		
Accession No. Uniprot GeneID Excitation Emission	Swiss-Prot#:P42261 P42261 2890; AF350: 346nm/442nm		
Accession No. Uniprot GeneID Excitation Emission	Swiss-Prot#:P42261 P42261 2890; AF350: 346nm/442nm AF405: 401nm/421nm		
Accession No. Uniprot GeneID Excitation Emission	Swiss-Prot#:P42261 P42261 2890; AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm		
Accession No. Uniprot GeneID Excitation Emission	Swiss-Prot#:P42261 P42261 2890; AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm		
Accession No. Uniprot GeneID Excitation Emission	Swiss-Prot#:P42261 P42261 2890; AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm		
Accession No. Uniprot GeneID Excitation Emission	Swiss-Prot#:P42261 P42261 2890; AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm		
Accession No. Uniprot GeneID Excitation Emission	Swiss-Prot#:P42261 P42261 2890; AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm		
Accession No. Uniprot GeneID Excitation Emission	Swiss-Prot#:P42261 P42261 2890; AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm		
Accession No. Uniprot GeneID Excitation Emission	Swiss-Prot#:P42261 P42261 2890; AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm 92 kDa		
Accession No. Uniprot GeneID Excitation Emission	Swiss-Prot#:P42261 P42261 2890; AF355: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm 92 kDa 0.011M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide		

Application Details	
Suggested Dilution:	
AF350 conjugated: most appl	lications: 1: 50 - 1:
AF405 conjugated: most appl	olications: 1: 50 - 1: 25
AF488 conjugated: most appl	olications: 1: 50 - 1: 25

AF555 conjugated: most applications: 1: 50 - 1: 250	
AF594 conjugated: most applications: 1: 50 - 1: 250	
AF647 conjugated: most applications: 1: 50 - 1: 250	
AF680 conjugated: most applications: 1: 50 - 1: 250	
AF750 conjugated: most applications: 1: 50 - 1: 250	

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Glutamate receptors mediate most excitatory neurotransmission in the brain and play an important role in neural plasticity, neural development and neurodegeneration. Ionotropic glutamate receptors are categorized into NMDA receptors and kainate/AMPA receptors, both of which contain glutamate-gated, cation-specific ion channels. Kainate/AMPA receptors are co-localized with NMDA receptors in many synapses and consist of seven structurally related subunits designated GluR-1 to -7. The kainate/AMPA receptors are primarily responsible for the fast excitatory neuro-transmission by glutamate whereas the NMDA receptors are functionally characterized by a slow kinetic and a high permeability for Ca2+ ions. The NMDA receptors consist of five subunits: epsilion 1, 2, 3, 4 and one zeta subunit. The zeta subunit is expressed throughout the brainstem whereas the four epsilon subunits display limited distribution.

Note: This product is for in vitro research use only