

CDC40 Conjugated Antibody

Catalog No: #C49200



Package Size: #C49200-AF350 100ul #C49200-AF405 100ul #C49200-AF488 100ul
 #C49200-AF555 100ul #C49200-AF594 100ul #C49200-AF647 100ul
 #C49200-AF680 100ul #C49200-AF750 100ul #C49200-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	CDC40 Conjugated Antibody
Host Species	Rabbit
Clonality	Monoclonal
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	Cdc40 antibody Cell division cycle 40 antibody Cell division cycle 40 homolog antibody EH binding protein 3 antibody EH-binding protein 3 antibody Ehb3 antibody hPRP17 antibody Pre mRNA processing factor 17 antibody Pre mRNA splicing factor 17 antibody Pre-mRNA-processing factor 17 antibody PRP17 antibody PRP17 homolog antibody PRP17_HUMAN antibody PRPF17 antibody
Accession No.	Swiss-Prot#:O60508
Uniprot	O60508
GeneID	51362;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	66 kDa
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Cell cycle events are regulated by the sequential activation and deactivation of cyclin dependent kinases (Cdks) and by the proteolysis of cyclins. The cell division control (Cdc) genes are required at various points in the cell cycle. Cdc40, also known as pre-mRNA-processing factor 17 (PRPF17) or EH-binding protein 3 (EHB3), is a 579 amino acid nuclear protein. Cdc40 is essential for the catalytic step II of the pre-mRNA splicing process, in which Cdc40 associates with the spliceosome C complex. Cdc40 contains seven WD repeats, which are important in protein-protein interactions. Cdc40 has sequence similarity to the yeast protein Prp17, which is involved in pre-mRNA splicing and cell cycle progression. The sequence similarity between the mammalian Cdc40 and the yeast Prp17 may indicate an additional role in cell cycle progression for mammalian Cdc40.

Note: This product is for in vitro research use only