FOXA1 Conjugated Antibody

Catalog No: #C49403

SAB Signalway Antibody

Package Size: #C49403-AF350 100ul #C49403-AF405 100ul #C49403-AF488 100ul

#C49403-AF555 100ul #C49403-AF594 100ul #C49403-AF647 100ul

#C49403-AF680 100ul #C49403-AF750 100ul #C49403-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	FOXA1 Conjugated Antibody
Host Species	Rabbit
Clonality	Monoclonal
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	forkhead box A1 antibody Forkhead box protein A1 antibody FOX A1 antibody FOXA1 antibody
	FOXA1_HUMAN antibody hepatocyte nuclear factor 3 alpha antibody Hepatocyte nuclear factor 3-alpha
	antibody HNF 3A antibody HNF-3-alpha antibody HNF-3A antibody HNF3A antibody MGC33105 antibody
	TCF 3A antibody TCF-3A antibody TCF3A antibody Transcription factor 3A antibody
Accession No.	Swiss-Prot#:P55317
Uniprot	P55317
GeneID	3169;
Excitation Emission	AF350: 346nm/442nm
	AF405: 401nm/421nm
	AF488: 493nm/519nm
	AF555: 555nm/565nm
	AF594: 591nm/614nm
	AF647: 651nm/667nm
	AF680: 679nm/702nm
	AF750: 749nm/775nm
Calculated MW	49 kDa
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
AF405 conjugated: most applications: 1: 50 - 1: 250
AF488 conjugated: most applications: 1: 50 - 1: 250
AF555 conjugated: most applications: 1: 50 - 1: 250
AF594 conjugated: most applications: 1: 50 - 1: 250
AF647 conjugated: most applications: 1: 50 - 1: 250
AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

HNF-1 (α and β), HNF-3 (α , β and γ), HNF-4 (α and γ), and HNF-6 compose, in part, a homoeprotein family designated the hepatocyte nuclear factor family. The various HNF-1 isoforms regulate transcription of genes in the liver as well as in other tissues such as kidney, small intestine and thymus. HNF-3 α , HNF-3 β and HNF-3 γ regulate the transcription of numerous hepatocyte genes in adult liver. HNF-3 α and HNF-3 β have also been shown to be involved in gastrulation events such as body axis formation. HNF-4 α and HNF-4 γ have been shown to be important for early embryo development. HNF-4 α is expressed in liver, kidney, pancreas, small intestine, testis and colon; HNF-4 γ is expressed in each of these tissues except liver. HNF-6 has been shown to bind to the promoter of HNF-3 β , which indicates a potential role of HNF-6 in gut endoderm epithelial cell differentiation. Evidence suggests that HNF-6 may also be a transriptional activator for at least 22 other hepatocyte-enriched genes, including cytochrome P450 2C13 and α -1 antitrypsin.

Note: This product is for in vitro research use only