

## Cullin 4a Conjugated Antibody

Catalog No: #C49742



Package Size: #C49742-AF350 100ul #C49742-AF405 100ul #C49742-AF488 100ul  
 #C49742-AF555 100ul #C49742-AF594 100ul #C49742-AF647 100ul  
 #C49742-AF680 100ul #C49742-AF750 100ul #C49742-Biotin 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)  
 Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Cullin 4a Conjugated Antibody
Host Species	Rabbit
Clonality	Monoclonal
Species Reactivity	Hu
Immunogen Description	Recombinant protein
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	2810470J21Rik antibody AW495282 antibody CUL 4A antibody CUL-4A antibody Cul4a antibody Cul4a protein antibody CUL4A_HUMAN antibody Cullin-4A antibody MGC36573 antibody MGC64071 antibody
Accession No.	Swiss-Prot#:Q13619
Uniprot	Q13619
GeneID	8451;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	88 kDa
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

## Application Details

## Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250  
 AF405 conjugated: most applications: 1: 50 - 1: 250  
 AF488 conjugated: most applications: 1: 50 - 1: 250  
 AF555 conjugated: most applications: 1: 50 - 1: 250  
 AF594 conjugated: most applications: 1: 50 - 1: 250  
 AF647 conjugated: most applications: 1: 50 - 1: 250  
 AF680 conjugated: most applications: 1: 50 - 1: 250  
 AF750 conjugated: most applications: 1: 50 - 1: 250

## Background

---

Core component of multiple cullin-RING-based E3 ubiquitin-protein ligase complexes which mediate the ubiquitination of target proteins. As a scaffold protein may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme. The E3 ubiquitin-protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1. The functional specificity of the E3 ubiquitin-protein ligase complex depends on the variable substrate recognition component. DCX(DET1-COP1) directs ubiquitination of JUN. DCX(DDB2) directs ubiquitination of XPC. DCX(DDB2) ubiquitinates histones H3-H4 and is required for efficient histone deposition during replication-coupled (H3.1) and replication-independent (H3.3) nucleosome assembly, probably by facilitating the transfer of H3 from ASF1A/ASF1B to other chaperones involved in histone deposition.

---

Note: This product is for in vitro research use only