## p150 CAF1 Conjugated Antibody

Catalog No: #C49950

olgilaliv

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Package Size: #C49950-AF350 100ul #C49950-AF405 100ul #C49950-AF488 100ul

#C49950-AF555 100ul #C49950-AF594 100ul #C49950-AF647 100ul

#C49950-AF680 100ul #C49950-AF750 100ul #C49950-Biotin 100ul

## Description

Product Name	p150 CAF1 Conjugated Antibody
Host Species	Rabbit
Clonality	Monoclonal
Species Reactivity	Hu
Immunogen Description	Recombinant protein within human p150 CAF1 aa 100-300.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	CAF 1 antibody CAF 1 subunit A antibody CAF antibody CAF I 150 kDa subunit antibody CAF I p150 antibody CAF-I subunit A antibody CAF-I 150 kDa subunit antibody CAF-I p150 antibody CAF1 antibody CAF1A_HUMAN antibody CAF1B antibody CAF1P150 antibody CHAF1A antibody Chromatin assembly factor 1 subunit A antibody Chromatin Assembly Factor 1 Subunit A p150 antibody Chromatin assembly factor I (150 kDa) antibody Chromatin assembly factor I p150 subunit antibody DCAF1 antibody hp150 antibody MGC71229 antibody P150 antibody
Accession No.	Swiss-Prot#:Q13111
Uniprot	Q13111
GeneID	10036;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	107 kDa (Predicted band size)
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

## **Application Details**

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
AF405 conjugated: most applications: 1: 50 - 1: 250
AF488 conjugated: most applications: 1: 50 - 1: 250
AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250
AF647 conjugated: most applications: 1: 50 - 1: 250
AF680 conjugated: most applications: 1: 50 - 1: 250
AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Background

Chromatin assembly factor-1 (CAF-1) is a multisubunit protein complex that comprises three polypeptide subunits known as p150, p60, and p48. CAF-1 is a nucleosome assembly factor that deposits newly synthesized and acetylated Histones H3/H4 into nascent chromatin during DNA replication. The p150 subunit of CAF-1 also supports the maintenance of heterochromatin, which requires the synthesis of both new histones and heterochromatin proteins and their orderly assembly during DNA replication. Heterochromatin is characterized as densely coiled chromatin that generally replicates late during S phase, has a low gene density, and contains large blocks of repetitive DNA that is relatively inaccessible to DNA-modifying reagents. In late S phase, p150 directly associates with heterochromatin associated proteins 1 (HP1 $\alpha$ , HP1 $\beta$  and HP1 $\gamma$ ). As cells prepare for mitosis, CAF-1 p150 and some HP1 progressively dissociate from heterochromatin, coinciding with the phosphorylation of histone H3. The HP1 proteins reassociate with chromatin at the end of mitosis, as histone H3 is dephosphorylated.

Note: This product is for in vitro research use only