Histone H4 (tri methyl K20) Conjugated Antibody

Catalog No: #CHW226



Package Size: #CHW226-AF350 100ul #CHW226-AF405 100ul #CHW226-AF488 100ul

#CHW226-AF555 100ul #CHW226-AF594 100ul #CHW226-AF647 100ul

#CHW226-AF680 100ul #CHW226-AF750 100ul #CHW226-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Storage	Store at + O in dark for 0 months
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide Store at 4°C in dark for 6 months
Calculated MW	60
2	AF750: 749nm/775nm
	AF680: 679nm/702nm
	AF647: 651nm/667nm
	AF594: 591nm/614nm
	AF555: 555nm/565nm
	AF488: 493nm/519nm
	AF405: 401nm/421nm
Excitation Emission	AF350: 346nm/442nm
GeneID	4089;
Uniprot	Q13485
Accession No.	Swiss-Prot#:Q13485
	against DPP homolog 4 antibody SMAD4 antibody SMAD4_HUMAN antibody
	SMA- and MAD-related protein 4 antibody SMAD 4 antibody SMAD family member 4 antibody SMAD mothers
	of, 4 antibody Mothers against DPP homolog 4 antibody MYHRS antibody OTTHUMP00000163548 antibody
	Mothers against decapentaplegic homolog 4 antibody Mothers against decapentaplegic, Drosophila, homolog
	decapentaplegic homolog 4 antibody MADH 4 antibody MADH4 antibody Medea antibody
	antibody MAD mothers against decapentaplegic Drosophila homolog 4 antibody MAD mothers against
	carcinoma 4 antibody DPC 4 antibody DPC4 antibody hSMAD4 antibody JIP antibody MAD homolog 4
	Pancreatic Carcinoma antibody Deleted in pancreatic carcinoma locus 4 antibody Deletion target in pancreati
Other Names	(Small) Mothers Against Decapentaplegic antibody Deleted in Pancreatic Carcinoma 4 antibody Deleted in
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Immunogen Description	recombinant protein
Species Reactivity	Hu, Ms, Rt
Host Species Clonality	Rabbit Monoclonal
Lloat Chasins	Dakkit

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250 AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250
AF555 conjugated: most applications: 1: 50 - 1: 250
AF594 conjugated: most applications: 1: 50 - 1: 250
AF647 conjugated: most applications: 1: 50 - 1: 250
AF680 conjugated: most applications: 1: 50 - 1: 250
AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Smad proteins, the mammalian homologs of the Drosophila Mothers against dpp (Mad) have been implicated as downstream effectors of TGF β /BMP signaling. Smad1 (also designated Madr1 or JV4-1), Smad5 and mammalian Smad8 (also designated Smad9 or MADH6) are effectors of BMP2 and BMP4 function while Smad2 (also designated Madr2 or JV18-1) and Smad3 are involved in TGF β and activin-mediated growth modulation. Smad4 (also designated DPC4) has been shown to mediate all of the above activities through interaction with various Smad family members. Smad6 and Smad7 regulate the response to activin/TGF β signaling by interfering with TGF β -mediated phosphorylation of other Smad family members.

Note: This product is for in vitro research use only