

H3F3B Conjugated Antibody

Catalog No: #C27332



Package Size: #C27332-AF350 100ul #C27332-AF405 100ul #C27332-AF488 100ul
 #C27332-AF555 100ul #C27332-AF594 100ul #C27332-AF647 100ul
 #C27332-AF680 100ul #C27332-AF750 100ul #C27332-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

| | |
|-----------------------|--|
| Product Name | H3F3B Conjugated Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Purification | Affinity purification |
| Applications | most applications |
| Species Reactivity | Hu |
| Immunogen Description | Recombinant fusion protein of human H3F3B (NP_002098.1). |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | H3F3B; H3.3B; histone H3.3 |
| Accession No. | Swiss-Prot#:P84243NCBI Gene ID:3021 |
| Uniprot | P84243 |
| GeneID | 3020;3021; |
| Excitation Emission | AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm |
| Calculated MW | 15kDa |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene contains introns and its mRNA is polyadenylated, unlike most histone genes. The protein encoded by this gene is a replication-independent histone that is a member of the histone H3 family. Pseudogenes of this gene have been identified on the X chromosome, and on chromosomes 5, 13 and 17.

Note: This product is for in vitro research use only