FBLIM1 Conjugated Antibody

Catalog No: #C29453



 Package Size:
 #C29453-AF350 100ul
 #C29453-AF405 100ul
 #C29453-AF488 100ul

 #C29453-AF555 100ul
 #C29453-AF594 100ul
 #C29453-AF647 100ul

 #C29453-AF680 100ul
 #C29453-AF750 100ul
 #C29453-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	FBLIM1 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Affinity purification
Applications	most applications
Species Reactivity	Hu,Rt
Immunogen Description	Recombinant fusion protein of human FBLIM1 (NP_060026.2).
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	FBLIM1; CAL; FBLP-1; FBLP1; filamin-binding LIM protein 1
Accession No.	Swiss-Prot#:Q8WUP2NCBI Gene ID:54751
Uniprot	Q8WUP2
GeneID	54751;
Excitation Emission	AF350: 346nm/442nm
	AF405: 401nm/421nm
	AF488: 493nm/519nm
	AF555: 555nm/565nm
	AF594: 591nm/614nm
	AF647: 651nm/667nm
	AF680: 679nm/702nm
	AF750: 749nm/775nm
Calculated MW	41kDa
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:	
AF350 conjugated: most applications: 1: 50 - 1: 250	
AF405 conjugated: most applications: 1: 50 - 1: 250	
AF488 conjugated: most applications: 1: 50 - 1: 250	
AF555 conjugated: most applications: 1: 50 - 1: 250	
AF594 conjugated: most applications: 1: 50 - 1: 250	
AF647 conjugated: most applications: 1: 50 - 1: 250	
AF680 conjugated: most applications: 1: 50 - 1: 250	

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

This gene encodes a protein with an N-terminal filamin-binding domain, a central proline-rich domain, and, multiple C-terminal LIM domains. This protein localizes at cell junctions and may link cell adhesion structures to the actin cytoskeleton. This protein may be involved in the assembly and stabilization of actin-filaments and likely plays a role in modulating cell adhesion, cell morphology and cell motility. This protein also localizes to the nucleus and may affect cardiomyocyte differentiation after binding with the CSX/NKX2-5 transcription factor. Alternative splicing results in multiple transcript variants encoding different isoforms.

Note: This product is for in vitro research use only