# NDUFA11 Conjugated Antibody

Catalog No: #C29788

Package Size: #C29788-AF350 100ul #C29788-AF405 100ul #C29788-AF488 100ul #C29788-AF555 100ul #C29788-AF594 100ul #C29788-AF647 100ul #C29788-AF680 100ul #C29788-AF750 100ul #C29788-Biotin 100ul

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com

## Description

| | |
|---|---|
| Product Name | NDUFA11 Conjugated Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Purification | Affinity purification |
| Applications | most applications |
| Species Reactivity | Ms |
| Immunogen Description | A synthetic peptide of human NDUFA11 (NP_783313.1). |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | NDUFA11; B14.7; CI-B14.7; NADH:ubiquinone oxidoreductase subunit A11 |
| Accession No. | Swiss-Prot#:Q86Y39NCBI Gene ID:126328 |
| Uniprot | Q86Y39 |
| GeneID | 126328; |
| Excitation Emission | AF350: 346nm/442nm<br>AF405: 401nm/421nm<br>AF488: 493nm/519nm<br>AF555: 555nm/565nm<br>AF594: 591nm/614nm<br>AF647: 651nm/667nm<br>AF680: 679nm/702nm<br>AF750: 749nm/775nm |
| Calculated MW | 15kDa |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

## Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Background

This gene encodes a subunit of the membrane-bound mitochondrial complex I. Complex I is composed of numerous subunits and functions as the NADH-ubiquinol reductase of the mitochondrial electron transport chain. Mutations in this gene are associated with severe mitochondrial complex I deficiency. Alternate splicing results in multiple transcript variants.

Note: This product is for in vitro research use only