FCN2 Conjugated Antibody

Catalog No: #C29937



 Package Size:
 #C29937-AF350 100ul
 #C29937-AF405 100ul
 #C29937-AF488 100ul

 #C29937-AF555 100ul
 #C29937-AF594 100ul
 #C29937-AF647 100ul

 #C29937-AF680 100ul
 #C29937-AF750 100ul
 #C29937-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	FCN2 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Isotype	lgG
Purification	Affinity purification
Applications	most applications
Species Reactivity	Hu,Ms,Rt
Immunogen Description	Recombinant fusion protein of human FCN2 (NP_004099.2).
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	FCN2; EBP-37; FCNL; P35; ficolin-2
Accession No.	Swiss-Prot#:Q15485NCBI Gene ID:2220
Uniprot	Q15485
GeneID	2220;
Excitation Emission	AF350: 346nm/442nm
	AF405: 401nm/421nm
	AF488: 493nm/519nm
	AF555: 555nm/565nm
	AF594: 591nm/614nm
	AF647: 651nm/667nm
	AF680: 679nm/702nm
	AF750: 749nm/775nm
Calculated MW	34kDa
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Formulation Storage	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:		
AF350 conjugated: most applications: 1: 50 - 1: 250		
AF405 conjugated: most applications: 1: 50 - 1: 250		
AF488 conjugated: most applications: 1: 50 - 1: 250		
AF555 conjugated: most applications: 1: 50 - 1: 250		
AF594 conjugated: most applications: 1: 50 - 1: 250		
AF647 conjugated: most applications: 1: 50 - 1: 250		
AF680 conjugated: most applications: 1: 50 - 1: 250		

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

The product of this gene belongs to the ficolin family of proteins. This family is characterized by the presence of a leader peptide, a short N-terminal segment, followed by a collagen-like region, and a C-terminal fibrinogen-like domain. This gene is predominantly expressed in the liver, and has been shown to have carbohydrate binding and opsonic activities. Alternatively spliced transcript variants encoding different isoforms have been identified.

Note: This product is for in vitro research use only