# DCDC2 Conjugated Antibody

Catalog No: #C30136

Package Size: #C30136-AF350 100ul #C30136-AF405 100ul #C30136-AF488 100ul #C30136-AF555 100ul #C30136-AF594 100ul #C30136-AF647 100ul #C30136-AF680 100ul #C30136-AF750 100ul #C30136-Biotin 100ul

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com

## Description

| | |
|---|---|
| Product Name | DCDC2 Conjugated Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Purification | Affinity purification |
| Applications | most applications |
| Species Reactivity | Rt |
| Immunogen Description | Recombinant fusion protein of human DCDC2 (NP_001182539). |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | DCDC2A, DFNB66, NPHP19, RU2, RU2S |
| Accession No. | Swiss-Prot#:Q9UHG0NCBI Gene ID:51473 |
| Uniprot | Q9UHG0 |
| GeneID | 51473; |
| Excitation Emission | AF350: 346nm/442nm<br>AF405: 401nm/421nm<br>AF488: 493nm/519nm<br>AF555: 555nm/565nm<br>AF594: 591nm/614nm<br>AF647: 651nm/667nm<br>AF680: 679nm/702nm<br>AF750: 749nm/775nm |
| Calculated MW | 53kDa |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

## Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Background

This gene encodes a doublecortin domain-containing family member. The doublecortin domain has been demonstrated to bind tubulin and enhance microtubule polymerization. This family member is thought to function in neuronal migration where it may affect the signaling of primary cilia. Mutations in this gene have been associated with reading disability (RD) type 2, also referred to as developmental dyslexia. Alternatively spliced transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jan 2013]

Note: This product is for in vitro research use only