H2AFV Conjugated Antibody

Catalog No: #C30191



 Package Size:
 #C30191-AF350 100ul
 #C30191-AF405 100ul
 #C30191-AF488 100ul

 #C30191-AF555 100ul
 #C30191-AF594 100ul
 #C30191-AF647 100ul

 #C30191-AF680 100ul
 #C30191-AF750 100ul
 #C30191-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

| Product Name | H2AFV Conjugated Antibody |
|-----------------------|---|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Purification | Affinity purification |
| Applications | most applications |
| Species Reactivity | Hu,Ms,Rt |
| Immunogen Description | Recombinant fusion protein of human H2AFV (NP_036544.1). |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | H2A.Z-2, H2AV |
| Accession No. | Swiss-Prot#:Q71UI9NCBI Gene ID:94239 |
| Uniprot | Q71UI9 |
| GeneID | 94239; |
| Excitation Emission | AF350: 346nm/442nm |
| | AF405: 401nm/421nm |
| | AF488: 493nm/519nm |
| | AF555: 555nm/565nm |
| | AF594: 591nm/614nm |
| | AF647: 651nm/667nm |
| | AF680: 679nm/702nm |
| | AF750: 749nm/775nm |
| Calculated MW | 14kDa |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| | |
| Storage | Store at 4°C in dark for 6 months |

Application Details

| Suggested Dilution: | |
|---|--|
| AF350 conjugated: most applications: 1: 50 - 1: 250 | |
| AF405 conjugated: most applications: 1: 50 - 1: 250 | |
| AF488 conjugated: most applications: 1: 50 - 1: 250 | |
| AF555 conjugated: most applications: 1: 50 - 1: 250 | |
| AF594 conjugated: most applications: 1: 50 - 1: 250 | |
| AF647 conjugated: most applications: 1: 50 - 1: 250 | |
| AF680 conjugated: most applications: 1: 50 - 1: 250 | |

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent histone that is a member of the histone H2A family. Several transcript variants encoding different isoforms, have been identified for this gene. [provided by RefSeq, Oct 2015]

Note: This product is for in vitro research use only