

PTPRF Conjugated Antibody

Catalog No: #C30653



Package Size: #C30653-AF350 100ul #C30653-AF405 100ul #C30653-AF488 100ul
 #C30653-AF555 100ul #C30653-AF594 100ul #C30653-AF647 100ul
 #C30653-AF680 100ul #C30653-AF750 100ul #C30653-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	PTPRF Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Affinity purification
Applications	most applications
Species Reactivity	Hu,Ms,Rt
Immunogen Description	Recombinant fusion protein of human PTPRF (NP_569707.2).
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	PTPRF; BNAH2; LAR; receptor-type tyrosine-protein phosphatase F
Accession No.	Swiss-Prot#:P10586NCBI Gene ID:5792
Uniprot	P10586
GeneID	5792;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	150kDa
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
 AF405 conjugated: most applications: 1: 50 - 1: 250
 AF488 conjugated: most applications: 1: 50 - 1: 250
 AF555 conjugated: most applications: 1: 50 - 1: 250
 AF594 conjugated: most applications: 1: 50 - 1: 250
 AF647 conjugated: most applications: 1: 50 - 1: 250
 AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP possesses an extracellular region, a single transmembrane region, and two tandem intracytoplasmic catalytic domains, and thus represents a receptor-type PTP. The extracellular region contains three Ig-like domains, and nine non-Ig like domains similar to that of neural-cell adhesion molecule. This PTP was shown to function in the regulation of epithelial cell-cell contacts at adherents junctions, as well as in the control of beta-catenin signaling. An increased expression level of this protein was found in the insulin-responsive tissue of obese, insulin-resistant individuals, and may contribute to the pathogenesis of insulin resistance. Two alternatively spliced transcript variants of this gene, which encode distinct proteins, have been reported.

Note: This product is for in vitro research use only