## KCTD7 Polyclonal Antibody

Catalog No: #46896

Package Size: #46896-1 50ul #46896-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| Descri | nti | Λn  |
|--------|-----|-----|
| DESCH  | DU  | ווט |
|        |     |     |

| Product Name          | KCTD7 Polyclonal Antibody  |
|-----------------------|--|
| Host Species          | Rabbit   |
| Clonality             | Polyclonal   |
| Purification          | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific |
|                       | immunogen.   |
| Applications          | WB   |
| Species Reactivity    | Hu Ms Rt   |
| Specificity           | This antibody detects endogenous levels of KCTD7   |
| Immunogen Type        | protein  |
| Immunogen Description | Synthesized peptide derived from human KCTD7. at AA range: 181-230   |
| Target Name           | KCTD7  |
| Other Names           | BTB/POZ domain-containing protein KCTD7  |
| Accession No.         | Swiss-Prot:Q96MP8NCBI Gene ID:154881   |
| Uniprot               | Q96MP8   |
| GeneID                | 154881;  |
| Calculated MW         | 33 kDa   |
| Concentration         | 1 mg/ml  |
| Formulation           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                    |
| Storage               | Store at -20°C   |
|                       |  |

## **Application Details**

Western blotting: 1B£B′1500-1B£B′12000

## **Images**



Western Blot analysis of 1,mouse-brain 2,mouse-spleen cells using primary antibody diluted at 1:1000(°C overnight). Secondary antibodyB£B1Goat Anti-rabbit IgG IRDye 800( diluted at 1:5000, 25B'F£C, 1 hour)

Note: This product is for in vitro research use only