ERBB2IP Conjugated Antibody

Catalog No: #C31608

SAB Signalway Antibody

Package Size: #C31608-AF350 100ul #C31608-AF405 100ul #C31608-AF488 100ul

#C31608-AF555 100ul #C31608-AF594 100ul #C31608-AF647 100ul

#C31608-AF680 100ul #C31608-AF750 100ul #C31608-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

| Product Name | ERBB2IP Conjugated Antibody |
|-----------------------|---|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Purification | Affinity purification |
| Applications | most applications |
| Species Reactivity | Hu |
| Immunogen Description | Recombinant fusion protein of human ERBB2IP (NP_061165.1). |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | ERBIN; ERBB2IP; HEL-S-78; LAP2; erbin |
| Accession No. | Swiss-Prot#:Q96RT1NCBI Gene ID:55914 |
| Uniprot | Q96RT1 |
| GeneID | 55914; |
| Excitation Emission | AF350: 346nm/442nm |
| | AF405: 401nm/421nm |
| | AF488: 493nm/519nm |
| | AF555: 555nm/565nm |
| | AF594: 591nm/614nm |
| | AF647: 651nm/667nm |
| | AF680: 679nm/702nm |
| | AF750: 749nm/775nm |
| Calculated MW | 182kDa |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |
| | |

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250
AF405 conjugated: most applications: 1: 50 - 1: 250
AF488 conjugated: most applications: 1: 50 - 1: 250
AF555 conjugated: most applications: 1: 50 - 1: 250
AF594 conjugated: most applications: 1: 50 - 1: 250
AF647 conjugated: most applications: 1: 50 - 1: 250
AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

This gene is a member of the leucine-rich repeat and PDZ domain (LAP) family. The encoded protein contains 17 leucine-rich repeats and one PDZ domain. It binds to the unphosphorylated form of the ERBB2 protein and regulates ERBB2 function and localization. It has also been shown to affect the Ras signaling pathway by disrupting Ras-Raf interaction. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

Note: This product is for in vitro research use only