

SAV1 Conjugated Antibody

Catalog No: #C31794



Package Size: #C31794-AF350 100ul #C31794-AF405 100ul #C31794-AF488 100ul
 #C31794-AF555 100ul #C31794-AF594 100ul #C31794-AF647 100ul
 #C31794-AF680 100ul #C31794-AF750 100ul #C31794-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

| | |
|-----------------------|--|
| Product Name | SAV1 Conjugated Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Purification | Affinity purification |
| Applications | most applications |
| Species Reactivity | Hu |
| Immunogen Description | Recombinant fusion protein of human SAV1 (NP_068590.1). |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | SAV1; SAV; WW45; WWP4; protein salvador homolog 1 |
| Accession No. | Swiss-Prot#:Q9H4B6NCBI Gene ID:60485 |
| Uniprot | Q9H4B6 |
| GeneID | 60485; |
| Excitation Emission | AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm |
| Calculated MW | 45kDa |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

WW domain-containing proteins are found in all eukaryotes and play an important role in the regulation of a wide variety of cellular functions such as protein degradation, transcription, and RNA splicing. This gene encodes a protein with two WW domains, a SARAH domain, and a coiled-coil region and is ubiquitously expressed in adult tissues. This protein binds to MST1 (mammalian sterile 20-like kinase 1) and promotes MST1-induced apoptosis. It has also been shown to bind to HAX1 (hematopoietic cell-specific protein 1 (HS1)-associated protein X-1) and to attenuate the anti-apoptotic effects of HAX1. Studies in human and mouse suggest this gene acts as a tumor suppressor.

Note: This product is for in vitro research use only