

RNF128 Conjugated Antibody

Catalog No: #C31813



Package Size: #C31813-AF350 100ul #C31813-AF405 100ul #C31813-AF488 100ul
 #C31813-AF555 100ul #C31813-AF594 100ul #C31813-AF647 100ul
 #C31813-AF680 100ul #C31813-AF750 100ul #C31813-Biotin 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	RNF128 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antigen affinity purification
Species Reactivity	Hu, Ms
Immunogen Description	Fusion protein of human RNF128
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Target Name	RNF128
Other Names	GRAIL
Accession No.	Swiss-Prot#: Q9BRQ8NCBI Protein#: BC063404
Uniprot	Q9BRQ8
GeneID	84883;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at -20°C/1 year

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

The protein encoded by this gene is a type I transmembrane protein that localizes to the endocytic pathway. This protein contains a RING zinc-finger motif and has been shown to possess E3 ubiquitin ligase activity. Expression of this gene in retrovirally transduced T cell hybridoma significantly inhibits activation-induced IL2 and IL4 cytokine production. Induced expression of this gene was observed in anergic CD4(+) T cells, which suggested a role in the induction of anergic phenotype. Alternatively spliced transcript variants encoding distinct isoforms have been reported.

Note: This product is for in vitro research use only