

## DDX47 Conjugated Antibody

Catalog No: #C31842



Package Size: #C31842-AF350 100ul #C31842-AF405 100ul #C31842-AF488 100ul  
 #C31842-AF555 100ul #C31842-AF594 100ul #C31842-AF647 100ul  
 #C31842-AF680 100ul #C31842-AF750 100ul #C31842-Biotin 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	DDX47 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antigen affinity purification
Species Reactivity	Hu, Ms
Immunogen Description	Fusion protein of human DDX47
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Target Name	DDX47
Other Names	RRP3; E4-DBP; HQ0256; MSTP162
Accession No.	Swiss-Prot#: Q07001 NCBI Protein#: BC009379
Uniprot	Q07001
GeneID	1144;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at -20°C/1 year

## Application Details

## Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## Background

---

This gene encodes a member of the DEAD box protein family. DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure, such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. The protein encoded by this gene can shuttle between the nucleus and the cytoplasm, and has an RNA-independent ATPase activity. Two alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.

---

Note: This product is for in vitro research use only