

TMX3 Conjugated Antibody

Catalog No: #C31877



Package Size: #C31877-AF350 100ul #C31877-AF405 100ul #C31877-AF488 100ul
 #C31877-AF555 100ul #C31877-AF594 100ul #C31877-AF647 100ul
 #C31877-AF680 100ul #C31877-AF750 100ul #C31877-Biotin 100ul

Orders: order@signalwayantibody.com
 Support: tech@signalwayantibody.com

Description

Product Name	TMX3 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antigen affinity purification
Species Reactivity	Hu, Ms
Immunogen Description	Fusion protein of human TMX3
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Target Name	TMX3
Other Names	PDIA13; TXNDC10
Accession No.	Swiss-Prot#: Q9H0I2NCBI Protein#: BC032325
Uniprot	Q9H0I2
GeneID	84080;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at -20°C/1 year

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The canonical protein encoded by this gene has an N-terminal ER-signal sequence, a catalytically active thioredoxin domain, one transmembrane domain and a C-terminal ER-retention sequence. This gene is expressed in many tissues but has its highest expression in heart and skeletal muscle. It is expressed in the retinal neuroepithelium and lens epithelium in the developing murine eye and haploinsufficiency of this gene in humans and zebrafish is associated with microphthalmia. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Note: This product is for in vitro research use only