

p53(Ab-15) Conjugated Antibody

Catalog No: #C21085



Package Size: #C21085-AF350 100ul #C21085-AF405 100ul #C21085-AF488 100ul

#C21085-AF555 100ul #C21085-AF594 100ul #C21085-AF647 100ul

#C21085-AF680 100ul #C21085-AF750 100ul #C21085-Biotin 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	p53(Ab-15) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of total p53 protein.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa.13~17 (P-L-S-Q-E) derived from Human p53.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Target Name	p53
Other Names	Tumor suppressor p53; Phosphoprotein p53; Antigen NY-CO-13; TP53;
Accession No.	Swiss-Prot: P04637NCBI Protein: NP_000537.3
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Concentration	1.0mg/ml
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details

Predicted MW: 53kd

Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

Immunofluorescence: 1:100~1:200

Background

p53 is a nuclear protein which plays an essential role in the regulation of cell cycle specifically in the transition from G0 to G1. It is found in very low

levels in normal cells however in a variety of transformed cell lines in high amounts and believed to contribute to transformation and malignancy. The open reading frame of p53 is 393 amino acids long, with the central region (consisting of amino acids from about 100 to 300) containing the DNA-binding domain. This proteolysis-resistant core is flanked by a C-terminal end mediating oligomerization and an N-terminal end containing a strong transcription activation signal. p53 binds as a tetramer to a PBS (p53-Binding Site) and activates the expression of downstream genes that inhibit growth and/or invasion. p53 binds as a tetramer to a p53-binding site (PBS) and to activate the expression of adjacent genes that inhibit growth and/or invasion. Deletion of one or both p53 alleles reduces the expression of tetramers, resulting in decreased expression of the growth inhibitory genes

Lin T, et al. (2005) Nat Cell Biol; 7(2): 165-71.

Vega FM, et al. (2004) Mol Cell Biol; 24(23): 10366-80.

Li J, et al. (2004) J Biol Chem; 279(40): 41275-9.

Wang J, et al. (2004) J Biol Chem; 279(38): 39584-92.

Note: This product is for in vitro research use only