

Chk2 Antibody

Catalog No: #24130

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	Chk2 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Affinity chromatography purified via peptide column
Applications	ELISA WB ICC
Species Reactivity	Hu Ms Rt
Immunogen Type	Peptide
Immunogen Description	Raised against a synthetic peptide corresponding to amino acids near the amino terminus of human Chk2.
Target Name	Chk2
Accession No.	Swiss-Prot:O96017Gene ID:11200
Uniprot	O96017
GeneID	11200;
Concentration	1mg/ml
Formulation	Supplied in PBS containing 0.02% sodium azide.
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.


Application Details

Predicted MW: 60 kd


Images



Western blot analysis of Chk2 expression in K562 (A), Jurkat (B), and HL-60 (C) whole cell lysates with Chk2 antibody at 1 ug /ml.



Immunocytochemistry of Chk2 in Jurkat cells with Chk2 antibody at 1 µg/mL.



Immunofluorescence of Chk2 in Jurkat cells with Chk2 antibody at 5 µg/mL

Background

The p53 tumor-suppressor gene integrates numerous signals that control cell life and death. Several novel molecules involved in p53 signaling, including Chk2, p53R2, p53AIP1, Noxa, PIDD, and PID/MTA2, were recently discovered. The checkpoint kinase Chk2 is the mammalian homologue of yeast Cds1/Rad53. In response to DNA damage, the checkpoint kinase ATM phosphorylates and activates Chk2, which in turn directly phosphorylates and activates p53. Chk2 serves as ATM downstream effector to mediate activation of p53. Chk2 also phosphorylates and activates BRCA1, the product of a tumor suppressor gene that is mutated in breast and ovarian cancer.

Note: This product is for in vitro research use only