MAP1 Antibody

Catalog No: #24162




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	Support: tech@signalwayantibody.com
Product Name	MAP1 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Affinity chromatography purified via peptide column
Applications	ELISA WB
Species Reactivity	Hu Ms
Immunogen Type	Peptide
Immunogen Description	Raised against a 15 amino acid peptide from near the center of human MAP-1.
Target Name	MAP1
Accession No.	Swiss-Prot:Q96BY2Gene ID:64112
Uniprot	Q96BY2
GeneID	64112;
Concentration	1mg/ml
Formulation	Supplied in PBS containing 0.02% sodium azide.

Images

Storage



Western blot analysis of MAP-1 in EL4 cell lysate with MAP-1 antibody (IN) at (A) 1, (B) 2, and (C) 4 ug/mL.

Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated

freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Background

Apoptosis plays a major role in normal organism development, tissue homeostasis, and removal of damaged cells. Disruption of this process has been implicated in a variety of diseases such as cancer. Members of the Bcl-2 family are known to be critical regulators of this process. These proteins are characterized by the presence of several conserved motifs termed Bcl-2 homology (BH) domains. A related protein termed MAP-1 has recently been identified. This protein contains a BH3-like domain and induces caspase-dependent apoptosis in mammalian cells when overexpressed. It forms homodimers and associates with Bcl-2 family members such as Bax, Bcl-2, and Bcl-XL in vitro and in vivo. It has been suggested that MAP-1 associates with the tumor suppressor RASSF1A following death receptor activation, allowing a conformational change in Bax that leads to cellular apoptosis.

Note: This product is for in vitro research use only