

PAK4 Antibody

Catalog No: #24183


Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


Description

Product Name	PAK4 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Affinity chromatography purified via peptide column
Applications	ELISA WB IHC
Species Reactivity	Hu
Immunogen Type	Peptide
Immunogen Description	Raised against a 13 amino acid peptide from near the center of human PAK4.
Target Name	PAK4
Other Names	p21-activated kinase 4
Accession No.	Swiss-Prot:O96013Gene ID:10298
Uniprot	O96013
GeneID	10298;
Concentration	1mg/ml
Formulation	Supplied in PBS containing 0.02% sodium azide.
Storage	Can be stored at -20°C, stable for one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Images



Western blot analysis of PAK4 in SW480 lysate with PAK4 antibody at (A) 1 and (B) 2 ug/mL.



Immunohistochemistry of PAK4 in human colon tissue with PAK4 antibody at 10 ug/mL.

Background

The p21-activated kinases (PAKs) are serine-threonine kinases that bind to the active forms of Cdc42 and Rac. They are divided into two groups, the first of which include PAK1, 2 and 3, and can be activated by Cdc42/Rac binding. Group 1 PAKs contain an autoinhibitory domain whose activity is regulated by Cdc42/Rac binding. The group 1 PAKs are known to be involved in cellular processes such as gene transcription, apoptosis, and cell morphology and motility. Much less is known about the second group, which includes PAK4, 5 and 6. These proteins are not activated by Cdc42/Rac binding. PAK4 was initially identified as a novel effector of Cdc42Hs. Co-expression of PAK4 and Cdc42Hs resulted in induction of filopodia and actin polymerization, showing that it is involved in cytoskeletal reorganization. Other experiments have shown PAK4 to be essential for embryonic viability and proper neuronal development. PAK4 has also been implicated in anchorage-independent growth of tumor cells and is required for activation of several cancer pro-survival pathways.

Note: This product is for in vitro research use only