

TPOR (phospho-Tyr626) rabbit pAb

Catalog No: #13471

Package Size: #13471-1 50ul #13471-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	TPOR (phospho-Tyr626) rabbit pAb
Host Species	Rabbit
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Applications	WB
Species Reactivity	Human
Specificity	This antibody detects endogenous levels of Human TPOR (phospho-Tyr626)
Immunogen Description	Synthesized phospho peptide around human TPOR (Tyr626)
Other Names	Thrombopoietin receptor (TPO-R) (Myeloproliferative leukemia protein) (Proto-oncogene c-Mpl) (CD antigen CD110)
Accession No.	Swiss Prot:P40238GenelD:4352
Uniprot	P40238
GeneID	4352
SDS-PAGE MW	69,40
Concentration	1 mg/ml
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	-20°C/1

Application Details

WB 1:1000-2000

Background

MPL proto-oncogene, thrombopoietin receptor(MPL) Homo sapiens In 1990 an oncogene, v-mpl, was identified from the murine myeloproliferative leukemia virus that was capable of immortalizing bone marrow hematopoietic cells from different lineages. In 1992 the human homologue, named, c-mpl, was cloned. Sequence data revealed that c-mpl encoded a protein that was homologous with members of the hematopoietic receptor superfamily. Presence of anti-sense oligodeoxynucleotides of c-mpl inhibited megakaryocyte colony formation. The ligand for c-mpl, thrombopoietin, was cloned in 1994. Thrombopoietin was shown to be the major regulator of megakaryocytopoiesis and platelet formation. The protein encoded by the c-mpl gene, CD110, is a 635 amino acid transmembrane domain, with two extracellular cytokine receptor domains and two intracellular cytokine receptor box motifs. TPO-R deficient mice were severely thrombocytopenic, emphasizing the important

Note: This product is for in vitro research use only