## PTEN (phospho Ser385) Polyclonal Antibody

Catalog No: #13595

Package Size: #13595-1 50ul #13595-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

## Description

| Beeenptien            |  |
|-----------------------|--|
| Product Name          | PTEN (phospho Ser385) Polyclonal Antibody  |
| Host Species          | Rabbit   |
| Purification          | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific |
|                       | immunogen.   |
| Applications          | IHC-p,IF(paraffin section),ELISA   |
| Species Reactivity    | Human,Mouse,Rat  |
| Specificity           | Phospho-PTEN (S385) Polyclonal Antibody detects endogenous levels of PTEN protein only when                |
|                       | phosphorylated at S385.  |
| Immunogen Description | The antiserum was produced against synthesized peptide derived from human PTEN around the                  |
|                       | phosphorylation site of Ser385. AA range:370-400   |
| Other Names           | PTEN; MMAC1; TEP1; Phosphatidylinositol 3; 4,5-trisphosphate 3-phosphatase and dual-specificity protein    |
|                       | phosphatase PTEN; Mutated in multiple advanced cancers 1; Phosphatase and tensin homolog                   |
| Accession No.         | Swiss Prot:P60484GeneID:5728   |
| Uniprot               | P60484   |
| GenelD                | 5728   |
| Concentration         | 1 mg/ml  |
| Formulation           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                    |
| Storage               | -20°C/1  |
|                       |  |

## **Application Details**

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

## Background

phosphatase and tensin homolog(PTEN) Homo sapiens This gene was identified as a tumor suppressor that is mutated in a large number of cancers at high frequency. The protein encoded by this gene is a phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase. It contains a tensin like domain as well as a catalytic domain similar to that of the dual specificity protein tyrosine phosphatases. Unlike most of the protein tyrosine phosphatases, this protein preferentially dephosphorylates phosphoinositide substrates. It negatively regulates intracellular levels of phosphatidylinositol-3,4,5-trisphosphate in cells and functions as a tumor suppressor by negatively regulating AKT/PKB signaling pathway. The use of a non-canonical (CUG) upstream initiation site produces a longer isoform that initiates translation with a leucine, and is thought to be preferentially associated with the mitochondrial inner membrane. This longer isoform may help regulate ener

Note: This product is for in vitro research use only