MCM4 (phospho Ser54) Polyclonal Antibody

Catalog No: #13725

Package Size: #13725-1 50ul #13725-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Decemption	
Product Name	MCM4 (phospho Ser54) Polyclonal Antibody
Host Species	Rabbit
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific
	immunogen.
Applications	WB,IHC-p,IF(paraffin section),ELISA
Species Reactivity	Human,Mouse,Monkey
Specificity	Phospho-MCM4 (S54) Polyclonal Antibody detects endogenous levels of MCM4 protein only when
	phosphorylated at S54.
Immunogen Description	The antiserum was produced against synthesized peptide derived from human MCM4 around the
	phosphorylation site of Ser54. AA range:20-69
Other Names	MCM4; CDC21; DNA replication licensing factor MCM4; CDC21 homolog; P1-CDC21
Accession No.	Swiss Prot:P33991GeneID:4173
Uniprot	P33991
GenelD	4173
SDS-PAGE MW	85
Concentration	1 mg/ml
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	-20°C/1

Application Details

Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.

Background

minichromosome maintenance complex component 4(MCM4) Homo sapiens The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance proteins (MCM) that are essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by MCM proteins is a key component of the pre-replication complex (pre_RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 6 and 7 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. The phosphorylation of this protein by CDC2 kinase reduces the DNA helicase activity and chromatin binding of the MCM complex. This gene is mapped to a region on the chromosome 8 head-to-head next to the PRKDC/DNA-PK, a DNA-activated protein kinase involved in the repair of DNA double-strand breaks. Alternatively spliced transcri

Note: This product is for in vitro research use only