

hnRNP D0 (phospho Ser83) Polyclonal Antibody

Catalog No: #13819



Package Size: #13819-1 50ul #13819-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

| | |
|-----------------------|---|
| Product Name | hnRNP D0 (phospho Ser83) Polyclonal Antibody |
| Host Species | Rabbit |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Applications | WB,IHC-p,IF/ICC,ELISA |
| Species Reactivity | Human,Mouse,Rat |
| Specificity | Phospho-hnRNP D0 (S83) Polyclonal Antibody detects endogenous levels of hnRNP D0 protein only when phosphorylated at S83. |
| Immunogen Description | The antiserum was produced against synthesized peptide derived from human hnRNP D0 around the phosphorylation site of Ser83. AA range:49-98 |
| Other Names | HNRNPD; AUF1; HNRPD; Heterogeneous nuclear ribonucleoprotein D0; hnRNP D0; AU-rich element RNA-binding protein 1 |
| Accession No. | Swiss Prot:Q14103GeneID:3184 |
| Uniprot | Q14103 |
| GeneID | 3184 |
| SDS-PAGE MW | 38 |
| Concentration | 1 mg/ml |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Storage | -20°C/1 |

Application Details

Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

Background

heterogeneous nuclear ribonucleoprotein D(HNRNPD) Homo sapiens This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are nucleic acid binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has two repeats of quasi-RRM domains that bind to RNAs. It localizes to both the nucleus and the cytoplasm. This protein is implicated in the regulation of mRNA stability. Alternative splicing of this gene results in four transcript variants. [provided by RefSeq, Jul 2008],

Note: This product is for in vitro research use only