## Flk-1 (phospho Tyr951) Polyclonal Antibody

Catalog No: #13864

Package Size: #13864-1 50ul #13864-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

## Description

Decomption	
Product Name	Flk-1 (phospho Tyr951) Polyclonal Antibody
Host Species	Rabbit
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific
	immunogen.
Applications	WB,IHC-p,IF/ICC,ELISA
Species Reactivity	Human,Mouse
Specificity	Phospho-Flk-1 (Y951) Polyclonal Antibody detects endogenous levels of Flk-1 protein only when
	phosphorylated at Y951.
Immunogen Description	The antiserum was produced against synthesized peptide derived from human VEGFR2 around the
	phosphorylation site of Tyr951. AA range:917-966
Other Names	KDR; FLK1; VEGFR2; Vascular endothelial growth factor receptor 2; VEGFR-2; Fetal liver kinase 1; FLK-1;
	Kinase insert domain receptor; KDR; Protein-tyrosine kinase receptor flk-1; CD antigen CD309
Accession No.	Swiss Prot:P35968GeneID:3791
Uniprot	P35968
GenelD	3791
Calculated MW	151kd
Concentration	1 mg/ml
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	-20°C/1

## Application Details

Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

## Background

kinase insert domain receptor(KDR) Homo sapiens Vascular endothelial growth factor (VEGF) is a major growth factor for endothelial cells. This gene encodes one of the two receptors of the VEGF. This receptor, known as kinase insert domain receptor, is a type III receptor tyrosine kinase. It functions as the main mediator of VEGF-induced endothelial proliferation, survival, migration, tubular morphogenesis and sprouting. The signalling and trafficking of this receptor are regulated by multiple factors, including Rab GTPase, P2Y purine nucleotide receptor, integrin alphaVbeta3, T-cell protein tyrosine phosphatase, etc.. Mutations of this gene are implicated in infantile capillary hemangiomas. [provided by RefSeq, May 2009],

Note: This product is for in vitro research use only