ATR (phospho-Thr1989) rabbit pAb

Catalog No: #14059

Package Size: #14059-1 50ul #14059-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Description | |
|-----------------------|-----------------------------------------------------------------------------------------------------------|
| Product Name | ATR (phospho-Thr1989) rabbit pAb |
| Host Species | Rabbit |
| Purification | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Applications | WB |
| Species Reactivity | Human |
| Specificity | This antibody detects endogenous levels of Human ATR (phospho-Thr1989) |
| Immunogen Description | Synthesized phosho peptide around human ATR (Thr1989) |
| Other Names | Serine/threonine-protein kinase ATR (EC 2.7.11.1) (Ataxia telangiectasia and Rad3-related protein) |
| | (FRAP-related protein 1) |
| Accession No. | Swiss Prot:Q13535GeneID:545 |
| Uniprot | Q13535 |
| GeneID | 545 |
| SDS-PAGE MW | 300 |
| Concentration | 1 mg/ml |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| | |

Application Details

WB 1:1000-2000

Storage

Background

ATR serine/threonine kinase(ATR) Homo sapiens The protein encoded by this gene belongs the PI3/PI4-kinase family, and is most closely related to ATM, a protein kinase encoded by the gene mutated in ataxia telangiectasia. This protein and ATM share similarity with Schizosaccharomyces pombe rad3, a cell cycle checkpoint gene required for cell cycle arrest and DNA damage repair in response to DNA damage. This kinase has been shown to phosphorylate checkpoint kinase CHK1, checkpoint proteins RAD17, and RAD9, as well as tumor suppressor protein BRCA1. Mutations of this gene are associated with Seckel syndrome. An alternatively spliced transcript variant of this gene has been reported, however, its full length nature is not known. Transcript variants utilizing alternative polyA sites exist. [provided by RefSeq, Jul 2008],

Note: This product is for in vitro research use only

-20°C/1