SARS-CoV-2 (2019-nCoV) S protein RBD, mFc Tag

Catalog No: #AP89618

SAB Signalway Antibody

Package Size: #AP89618-1 10ug #AP89618-2 100ug

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	SARS-CoV-2 (2019-nCoV) S protein RBD, mFc Tag
Host Species	HEK293
Purification	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Species Reactivity	Human
Immunogen Description	S protein RBD(Arg319-Phe541)+mFc(Pro99-Lys330)
Other Names	S protein RBD, Spike glycoprotein Receptor-binding domain, S glycoprotein RBD, Spike protein RBD,
	COVID-19
Calculated MW	51.3 kDa
Tag Info	C-Mouse Fc tag
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 98 % trehalose is added as protectants before
	lyophilization.
Storage	Store at -80°C for 12 months (Avoid repeated freezing and thawing)

Images




Figure 1. SARS-CoV-2 (2019-nCoV) S protein RBD, mFc Tag on SDS-PAGE under reducing condition.




Figure 2. ELISA plate pre-coated by 2 µg/ml (100 µl/well) S-RBD, mFc tagged protein can bind Human ACE2, hFc Tagged protein in a linear range of 7.81-87.7 ng/ml.

Product Description

Expression Region:620Research Topic:SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as Covid19 (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. The spike protein is a type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which accounts for recognizing the cell surface receptor, ACE2. S2 contains basic elements needed for the membrane fusion. Recent publications indicate that S1-RBD domain can induce virus neutralizing-antibody and T cell response.

Note: This product is for in vitro research use only