

## Recombinant human IL4

Catalog No: #AG0003

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Recombinant human IL4
Host Species	HEK293
Purification	> 95% by Tris-Bis PAGE;> 95% by SEC-HPLC
Immunogen Description	His25-Ser153
Target Name	IL4
Other Names	Human IL-4, h-IL-4, rh-IL-4, recombinant IL-4, interleukin-4
Accession No.	Uniprot:P05112Gene ID:3565
Uniprot	P05112
GeneID	3565
Target Species	human
Calculated MW	14.9 KDa
Tag Info	additional amino acid free
Formulation	0.22 µm filtered solution of PBS, pH 7.4.
Storage	Aliquot and store at -80°C. Avoid repeated freeze/thaw cycles.

## Background

Interleukin-4 (IL-4), also known as B cell-stimulatory factor-1, is a monomeric, approximately 13?kDa?18?kDa Th2 cytokine that shows pleiotropic effects during immune responses (1?3). It is a glycosylated polypeptide that contains three intrachain disulfide bridges and adopts a bundled four alpha -helix structure (4). Human IL-4 is synthesized with a 24 aa signal sequence. Alternate splicing generates an isoform with a 16 aa internal deletion. Mature human IL-4 shares 55%, 39% and 43% aa sequence identity with bovine, mouse, and rat IL-4, respectively. Human, mouse, and rat IL-4 are species-specific in their activities (5?7). IL-4 exerts its effects through two receptor complexes (8, 9). The type I receptor, which is expressed on hematopoietic cells, is a heterodimer of the ligand binding IL-4 R alpha and the common gamma ?chain (a shared subunit of the receptors for IL-2, -7, -9, -15, and ?21). The type II receptor on nonhematopoietic cells consists of IL-4 R alpha and IL?13?R alpha 1. The type II receptor also transduces IL-13 mediated signals. IL-4 is primarily expressed by Th2-biased CD4+?T cells, mast cells, basophils, and eosinophils (1, 2). It promotes cell proliferation, survival, and immunoglobulin class switch to IgG4 and IgE in human B cells, acquisition of the Th2 phenotype by na?ve CD4+?T cells, priming and chemotaxis of mast cells, eosinophils, and basophils, and the proliferation and activation of epithelial cells (10?13). IL-4 plays a dominant role in the development of allergic inflammation and asthma (12, 14).

Note: This product is for in vitro research use only