

## Recombinant mouse IL7

Catalog No: #AG0039

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Recombinant mouse IL7
Host Species	HEK293
Purification	> 95% by Tris-Bis PAGE;> 95% by SEC-HPLC
Immunogen Description	Glu26-Ile154
Target Name	IL7
Other Names	Mouse IL7; mIL-7; IL-7interleukin-7; interleukin 7; Lymphopoietin-1; PBGF
Accession No.	Uniprot:P10168Gene ID:16196
Uniprot	P10168
GeneID	16196
Target Species	mouse
Calculated MW	14.9 KDa
Tag Info	additional amino acid free
Formulation	0.22 µm filtered solution of PBS, pH 7.4.
Storage	Aliquot and store at -80°C. Avoid repeated freeze/thaw cycles.

## Background

IL-7 (interleukin-7) is a 25 kDa cytokine of the hemopoietin family that plays important roles in lymphocyte differentiation, proliferation, and survival (1-4). Mouse IL-7 cDNA encodes 154 amino acids (aa) that include a 25 aa signal peptide (4). Mouse IL-7 shares approximately 88% aa sequence identity with rat IL-7 and 58-60% with human, equine, bovine, ovine, porcine, feline and canine IL-7. Human and mouse IL-7 exhibit cross-species activity (2, 3). IL-7 is produced by a wide variety of cells in primary and secondary lymphoid tissues, including stromal epithelial cells of the thymus, bone marrow, and intestines (1, 2, 5). Circulating IL-7 is limiting in healthy animals, but increases during lymphopenia (1, 6). IL-7 signals through a complex of the IL-7 Receptor alpha subunit (IL-7 R alpha, also known as CD127) with the common gamma chain (gamma c) (1). The gamma c is also a subunit of the receptors for IL-2, -4, -9, -15, and -21 (1). IL-7 R alpha is expressed on double negative (CD4-CD8-) and CD4+ or CD8+ single positive na?ve and memory T cells, but undergoes IL-7-mediated down?regulation and shedding during antigen-driven T cell proliferation, and is absent on regulatory T cells (1, 2, 6-11). IL-7 contributes to the maintenance of all na?ve and memory T cells, mainly by promoting expression of the anti-apoptotic protein Bcl-2 (9-11). It is required for optimal T cell-dendritic cell interaction (6). IL-7 is expressed early in B cell development prior to the appearance of surface IgM (1, 5, 9). In mouse, IL-7 activation of IL-7 R alpha is critical for both T cell and B cell lineage development, while in humans, it is required for T cell but not for B cell development (4, 9, 12, 13). However, IL-7 functions in both mouse and human pro-B cells to suppress premature Ig light chain recombination during proliferative growth (14, 15).

Note: This product is for in vitro research use only