

# Recombinant human Trop2

Catalog No: #AG0063

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Recombinant human Trop2
Host Species	HEK293
Purification	> 95% by Tris-Bis PAGE; > 95% by SEC-HPLC
Immunogen Description	His27-Thr274
Target Name	Trop2
Other Names	Cell surface glycoprotein Trop-2; EGP1; EGP-1; epithelial glycoprotein-1; GA733-1; GA733-1GP50; gastrointestinal tumor-associated antigen GA7331; gp50; M1S1 cell surface glycoprotein TROP2
Accession No.	Uniprot:P09758 Gene ID:4070
Uniprot	P09758
GeneID	4070
Target Species	human
Calculated MW	28.7 kDa
Tag Info	C-His-Tag
Formulation	0.22 µm filtered solution of PBS, pH 7.4.
Storage	Aliquot and store at -80°C. Avoid repeated freeze/thaw cycles.

## Background

Human TROP-2, also called tumor associated calcium signal transducer 2 (TACSTD2), GA733-1, gp50 and T16, is a type I cell surface glycoprotein that is highly expressed on human carcinomas. It was originally identified as an antigen present on human gastrointestinal tumors and is the second of two members of this family. The other family member is GA733-2, also called EpCAM, TROP-1, 17-1A, gp40 and KSA. The TROP-2 gene is unique in that it contains no introns. A study of these two genes suggested that TROP-2 was the result of a retroposition of the EpCAM gene. TROP-2 and EpCAM share approximately 49% amino acid identity and approximately 67% similarity. Human and mouse TROP-2 share 87% similarity. The human TROP-2 protein consists of a putative 26 amino acid (aa) signal sequence, a 248 aa extracellular domain, a 23 aa transmembrane region and a 26 aa cytoplasmic domain. TROP-2 is capable of transducing an intracellular calcium signal and may play a role in tumor growth. It also has adhesive functions.

Note: This product is for in vitro research use only