

Recombinant human Spliceosome RNA helicase DDX39B



Catalog No: #AP71657

Orders: order@signalwayantibody.com

Package Size: #AP71657-1 20ug #AP71657-2 100ug

Support: tech@signalwayantibody.com

Description

Product Name	Recombinant human Spliceosome RNA helicase DDX39B
Brief Description	Recombinant Protein
Host Species	E.coli
Purification	Greater than 90% as determined by SDS-PAGE.
Immunogen Description	Expression Region:2-251aaSequence Info:Partial
Other Names	56KDA U2AF65-associated protein;ATP-dependent RNA helicase p47DEAD box protein UAP56HLA-B-associated transcript 1 protein
Accession No.	Q13838
Uniprot	Q13838
GeneID	7919;
Calculated MW	55.2 kDa
Tag Info	N-terminal GST-tagged
Target Sequence	AENDVDNELLDYEDDEVETAAGGDGAEAPAKKDVKGSYVSIHSSGFRDFFLLKPELLRAIVDCGFEHPSEVQHE CIPQAILGMDVLCQAKSGMGKTAVFVLTALQLEPVTGQVSVLVMCHTRELAFQISKEYERFSKYMPNVKVAV FFGGLSIKKDEEVLKKNCPHIVGTPGRILALARNKSLNLKHIKHFILDECDKMLEQLDMRRDVQEIFRMPHEK QVMFMSATLSKEIRPVCRKFMQDPMEIFV
Formulation	Tris-based buffer50% glycerol
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C,-80°C. The shelf life of lyophilized form is 12 months at -20°C,-80°C.Notes:Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

Background

Involved in nuclear export of spliced and unspliced mRNA. Assblng component of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and specifically associates with spliced mRNA and not with unspliced pre-mRNA. TREX is recruited to spliced mRNAs by a transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export to the cytoplasm via the TAP,NFX1 pathway. May undergo several rounds of ATP hydrolysis during assbly of TREX to drive subsequent loading of components such as ALYREF,THOC and CHTOP onto mRNA. The TREX complex is essential for the export of Kaposi's sarcoma-associated herpesvirus (KSHV) intronless mRNAs and infectious virus production. Also associates with pre-mRNA independent of ALYREF,THOC4 and the THO complex. Involved in the nuclear export of intronless mRNA; the ATP-bound form is proposed to recruit export adapter ALYREF,THOC4 to intronless mRNA; its ATPase activity is cooperatively stimulated by RNA and ALYREF,THOC4 and ATP hydrolysis is thought to trigger the dissociation from RNA to allow the association of ALYREF,THOC4 and the NXF1-NXT1 heterodimer. Involved in transcription elongation and genome stability.Splice factor that is required for the first ATP-dependent step in spliceosome assbly and for the interaction of U2 snRNP with the branchpoint. Has both RNA-stimulated ATP binding,hydrolysis activity and ATP-dependent RNA unwinding activity. Even with the stimulation of RNA, the ATPase activity is weak. Can only hydrolyze ATP but not other NTPs. The RNA stimulation of ATPase activity does not have a strong preference for the sequence and length of the RNA. However, ssRNA stimulates the

ATPase activity much more strongly than dsRNA. Can unwind 5' or 3' overhangs or blunt end RNA duplexes in vitro. The ATPase and helicase activities are not influenced by U2AF2; the effect of ALYREF,THOC4 is reported conflictingly with [PubMed:23299939] reporting a stimulatory effect.

References

The BAT1 gene in the MHC encodes an evolutionarily conserved putative nuclear RNA helicase of the DEAD family. Peelman L., Chardon P., Nunes M., Renard C., Geffrotin C., Vaiman M., van Zeveren A., Coppieters W., van de Weghe A., Bouquet Y., Choy W., Strominger J., Spies T. *Genomics* 26:210-218(1995) Research Topic: Epigenetics and Nuclear Signaling

Note: This product is for in vitro research use only