

Recombinant human Granulocyte colony-stimulating factor receptor

Catalog No: #AP72132

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Package Size: #AP72132-1 20ug #AP72132-2 100ug #AP72132-3 1mg

Description

Product Name	Recombinant human Granulocyte colony-stimulating factor receptor
Brief Description	Recombinant Protein
Host Species	E.coli
Purification	Greater than 90% as determined by SDS-PAGE.
Immunogen Description	Expression Region:25-620aaSequence Info:Extracellular Domain
Other Names	Colony-stimulating factor ;CSFMolgramostin;Sargramostim
Accession No.	Q99062
Uniprot	Q99062
GeneID	1441;
Calculated MW	93.5 kDa
Tag Info	N-terminal GST-tagged
Target Sequence	ECGHISVSAPIVHLGDPITASCIKQNCSHLDPEPQILWRLGAELQPGGRQQRLSDGTQESIITLPHLNHTQAF SCCLNWGNSLQILDQVELRAGYPPAIPHNLSCLMNLTSSLICQWEPGPETHLPTSFTLKSFKSRGNCQTQGD SILDCVPKDGQSHCCIPRKHLLLYQNMGIWVQAENALGTSMSPLCLDPMDDVVKLEPPMLRTMDPSPEAAPP QAGCLQLCWEPWQPLHINQKCELRHKPQRGEASWALVGPLLEALQYELCGLLPATAYTLQIRCIRWPLPG HWSDWSPSLELRTERAPTVRDLTWWRQRQLDPRTVQLFWKVPLEEDSGRIQGYVSWRPSGQAGAILPL CNTTELSCTFHLPSEAQEVAVAYNSAGTSRPTPVVFSES RGPALTRLHAMARDPHSLWVGWEPPNPWPQG YVIEWGLGPPSASNSNKTWRMEQNGRATGFLLENIRPFQLYEIIIVTPLYQDTMGPSQHVYAYSQEMAPSHA PELHLKHIGKTWAQLEWVPEPELGGKSPKTHYIFWTNAQNSFSAILNASSRGRFVHLGLEPASLYHIHLMAAS QAGATNSTVLTMLTLT
Formulation	Tris-based buffer50% glycerol
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C,-80°C. The shelf life of lyophilized form is 12 months at -20°C,-80°C.Notes:Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

Background

Cytokine that stimulates the growth and differentiation of hematopoietic precursor cells from various lineages, including granulocytes, macrophages, eosinophils and erythrocytes.

References

SeattleSNPs variation discovery resourceThe DNA sequence and comparative analysis of human chromosome 5.Schmutz J., Martin J., Terry A., Couronne O., Grimwood J., Lowry S., Gordon L.A., Scott D., Xie G., Huang W., Hellsten U., Tran-Gyamfi M., She X., Prabhakar S., Aerts A., Altherr M., Bajorek E., Black S., Branscomb E., Caoile C., Challacombe J.F., Chan Y.M., Denys M., Detter J.C., Escobar J., Flowers D., Fotopulos D.,

Glavina T., Gomez M., Gonzales E., Goodstein D., Grigoriev I., Groza M., Hammon N., Hawkins T., Haydu L., Israni S., Jett J., Kadner K., Kimball H., Kobayashi A., Lopez F., Lou Y., Martinez D., Medina C., Morgan J., Nandkeshwar R., Noonan J.P., Pitluck S., Pollard M., Predki P., Priest J., Ramirez L., Retterer J., Rodriguez A., Rogers S., Salamov A., Salazar A., Thayer N., Tice H., Tsai M., Ustaszewska A., Vo N., Wheeler J., Wu K., Yang J., Dickson M., Cheng J.-F., Eichler E.E., Olsen A., Pennacchio L.A., Rokhsar D.S., Richardson P., Lucas S.M., Myers R.M., Rubin E.M. Nature 431:268-274(2004) Research Topic: Cancer

Note: This product is for in vitro research use only